

Table 6B
2004 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Page: 1
Run Date: 10 / 11 / 05

Comparison of Growth and Anemia Indicators by County (3,4)													
County	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)						
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Alameda	4,112	7.4 (33)	10.7 (45)	22,579	5.0(34)	6.2(30)	14.2(29)	7,651	16.4(33)	16.4(23)		7,392	14.2 (34)
Alpine	32	* *	* *	164	8.5(64)	4.3(10)	15.2(37)	65	* *	* *		94	* *
Amador	157	10.8 (54)	5.7 (5)	737	5.2(41)	6.6(37)	14.9(35)	251	14.3(10)	12.7 (3)		329	11.6 (14)
Butte	1,806	7.7 (40)	7.9 (17)	6,833	5.2(41)	6.3(32)	12.0(13)	2,690	15.8(20)	13.5 (9)		4,510	10.1 (7)
Calaveras	159	10.7 (53)	8.2 (23)	988	6.7(60)	5.5(24)	18.8(60)	425	16.2(27)	16.5(24)		619	14.9 (38)
City Of Be	20	* *	* *	105	2.9 (1)	3.8 (3)	21.9(64)	27	* *	* *		61	* *
Colusa	239	6.7 (23)	5.0 (3)	1,245	3.6 (6)	8.2(51)	15.9(40)	486	13.2 (3)	22.2(57)		736	15.8 (45)
Contra Cos	2,755	7.0 (27)	10.6 (44)	13,961	4.7(30)	7.0(42)	14.5(30)	4,909	17.2(46)	17.4(32)		4,798	11.9 (16)
Del Norte	163	4.3 (3)	11.0 (48)	851	3.8 (8)	4.2 (8)	16.8(51)	342	19.6(58)	19.6(48)		567	7.8 (2)
El Dorado	516	7.4 (33)	6.4 (6)	2,011	5.1(37)	11.8(62)	9.9 (4)	682	14.1 (9)	13.2 (7)		1,007	10.4 (9)
Fresno	8,288	7.2 (29)	7.9 (17)	44,635	5.4(45)	4.0 (5)	16.5(47)	18,664	15.8(20)	17.0(30)		30,472	12.4 (18)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2004 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Page: 2
Run Date: 10 / 11 / 05

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low		High	Short Stature (6)			Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)			
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Glenn	210	4.3 (3)	17.6 (57)	1,096	5.0(34)	4.8(17)	16.1(42)	432	19.0(56)	19.2(46)	746	13.3 (27)
Humboldt	582	8.2 (45)	13.6 (56)	3,253	4.5(24)	5.6(26)	13.5(22)	1,165	18.5(52)	14.8(11)	1,855	12.5 (19)
Imperial	1,549	7.1 (28)	8.1 (22)	6,158	3.2 (4)	6.5(36)	15.3(38)	2,536	15.7(18)	18.0(37)	4,242	6.6 (1)
Inyo	91	* *	* *	475	5.5(46)	5.5(24)	18.1(54)	171	13.5 (4)	21.1(55)	262	15.6 (42)
Kern	6,678	7.2 (29)	8.8 (31)	32,396	4.5(24)	5.4(22)	16.1(42)	13,174	16.1(25)	16.8(29)	19,769	10.7 (11)
Kings	643	5.4 (9)	10.0 (41)	3,648	4.1(15)	4.8(17)	19.5(62)	1,453	16.4(33)	21.1(55)	2,361	27.5 (60)
L.A. East	16,856	6.9 (25)	7.9 (17)	43,777	6.0(51)	4.4(12)	18.1(54)	13,204	16.5(36)	17.6(34)	21,366	15.7 (43)
L.A. North	6,721	6.6 (20)	7.5 (12)	26,910	6.6(57)	4.6(14)	18.3(56)	8,601	17.3(47)	22.5(58)	12,586	13.3 (27)
L.A. Other	13,921	8.1 (44)	7.8 (14)	162,088	4.0(13)	4.3(10)	18.3(56)	81,396	16.4(33)	18.3(39)	105,642	13.6 (30)
L.A. South	16,511	8.2 (45)	7.6 (13)	45,834	6.2(55)	4.1 (7)	18.4(59)	14,055	16.3(31)	17.4(32)	22,945	13.8 (31)
L.A. West	3,685	6.2 (14)	6.6 (8)	10,925	6.1(53)	4.4(12)	17.6(52)	3,226	17.1(43)	18.5(40)	4,997	17.3 (53)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2004 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Page: 3
Run Date: 10 / 11 / 05

Comparison of Growth and Anemia Indicators by County (3,4)													
County	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)		Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Lake	215	9.3 (51)	8.8 (31)	1,012	3.0 (2)	5.4(22)	10.4 (6)	372	14.0 (7)	14.2(10)		666	13.1 (25)
Lassen	151	7.3 (32)	11.3 (51)	592	4.6(26)	9.5(58)	10.3 (5)	235	11.9 (1)	11.1 (2)		322	10.2 (8)
Long Beach	2,535	7.9 (42)	8.4 (26)	11,704	5.0(34)	5.1(21)	18.0(53)	4,810	16.6(39)	19.7(50)		5,999	13.3 (27)
Madera	1,582	6.5 (19)	8.5 (28)	5,617	4.6(26)	4.2 (8)	16.5(47)	2,313	17.7(50)	16.7(27)		4,302	16.5 (50)
Marin	110	4.5 (5)	5.5 (4)	2,446	3.8 (8)	8.2(51)	11.3 (8)	836	15.6(17)	18.5(40)		1,434	8.0 (3)
Mariposa	28	* *	* *	158	7.0(61)	12.0(64)	8.2 (2)	57	* *	* *		83	* *
Mendocino	699	8.0 (43)	10.0 (41)	2,666	3.2 (4)	8.6(55)	12.3(17)	972	18.8(53)	19.7(50)		1,041	13.9 (32)
Merced	2,567	7.6 (39)	8.5 (28)	10,723	5.8(50)	4.0 (5)	16.7(50)	4,177	18.2(51)	18.9(44)		7,964	15.5 (41)
Modoc	28	* *	* *	143	4.9(31)	6.3(32)	11.9(10)	63	* *	* *		91	* *
Mono	140	14.3 (57)	6.4 (6)	373	4.3(20)	11.8(62)	7.2 (1)	101	13.9 (5)	6.9 (1)		182	14.3 (35)
Monterey	1,069	5.1 (7)	10.9 (47)	8,743	5.5(46)	3.7 (1)	16.1(42)	2,667	19.4(57)	18.6(42)		4,666	12.5 (19)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2004 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Page: 4
Run Date: 10 / 11 / 05

Comparison of Growth and Anemia Indicators by County (3,4)												
County	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)	
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)					
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
Napa	462	7.4 (33)	9.1 (34)	2,593	4.0(13)	7.8(49)	12.2(16)	909	16.5(36)	15.3(15)	1,248	22.0 (59)
Nevada	365	5.8 (11)	7.1 (10)	1,253	3.8 (8)	10.3(60)	8.5 (3)	370	16.8(42)	12.7 (3)	580	12.8 (23)
Orange	10,888	6.4 (17)	8.0 (21)	29,231	7.3(62)	5.7(28)	18.3(56)	10,993	15.9(22)	19.6(48)	27,501	14.5 (37)
Pasadena	592	9.6 (52)	4.2 (1)	2,694	4.1(15)	6.6(37)	13.5(22)	1,075	14.0 (7)	13.0 (5)	1,841	8.9 (4)
Placer	686	6.3 (15)	8.6 (30)	2,824	6.1(53)	8.7(56)	11.4 (9)	918	13.9 (5)	15.8(18)	1,548	14.4 (36)
Plumas	84	* *	* *	375	5.1(37)	10.4(61)	14.7(33)	144	12.5 (2)	20.1(54)	186	11.8 (15)
Riverside	9,756	6.9 (25)	7.8 (14)	37,950	5.6(48)	4.7(16)	16.5(47)	13,018	15.1(11)	15.9(20)	19,170	16.0 (47)
Sacramento	3,748	7.2 (29)	9.8 (38)	36,353	5.1(37)	6.6(37)	13.9(26)	14,435	15.9(22)	15.5(16)	19,303	15.7 (43)
San Benito	176	4.5 (5)	11.4 (52)	1,332	4.6(26)	4.6(14)	19.8(63)	496	20.4(59)	25.4(59)	1,061	16.4 (49)
San Bernar	8,656	7.5 (36)	7.3 (11)	44,873	6.6(57)	4.9(20)	16.4(46)	16,384	15.9(22)	16.7(27)	23,944	16.7 (51)
San Diego	12,606	6.6 (20)	9.4 (36)	55,231	4.6(26)	6.2(30)	13.6(25)	20,629	15.7(18)	15.6(17)	35,302	12.5 (19)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2004 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Page: 5
Run Date: 10 / 11 / 05

Comparison of Growth and Anemia Indicators by County (3,4)													
County	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)						
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
San Franci	675	6.4 (17)	8.4 (26)	6,060	4.3(20)	7.2(45)	12.6(18)	2,049	16.7(41)	17.9(36)		1,044	11.3 (12)
San Joaqui	5,126	9.2 (50)	7.9 (17)	20,971	5.7(49)	6.0(29)	16.0(41)	8,042	16.2(27)	16.5(24)		8,485	11.4 (13)
San Luis O	1,129	8.3 (47)	11.2 (50)	4,468	4.1 (15)	9.6(59)	12.0(13)	1,484	16.2(27)	15.8(18)		1,975	16.8 (52)
San Mateo	930	6.8 (24)	9.1 (34)	10,653	4.4(22)	7.3(47)	14.5(30)	3,632	18.8(53)	19.7(50)		4,921	9.8 (6)
Santa Barb	3,596	14.2 (56)	7.0 (9)	14,162	4.9(31)	5.6(26)	14.9(35)	5,307	17.1(43)	19.7(50)		9,493	13.1 (25)
Santa Clar	2,650	5.3 (8)	9.4 (36)	27,303	4.1(15)	6.4(34)	13.3(20)	9,880	15.1(11)	16.5(24)		12,572	15.2 (40)
Santa Cruz	709	5.4 (9)	9.0 (33)	3,599	5.1(37)	7.0(42)	13.2(19)	1,039	15.1(11)	18.2(38)		1,482	14.9 (38)
Shasta	1,129	8.5 (48)	9.9 (39)	4,410	6.6(57)	3.7 (1)	14.7(33)	1,728	17.1(43)	15.0(13)		2,785	10.6 (10)
Sierra	13	* *	* *	53	* *	* *	* *	16	* *	* *		27	* *
Siskiyou	237	6.3 (15)	4.2 (1)	1,164	4.9(31)	9.1(57)	11.9(10)	465	17.4(48)	14.8(11)		807	18.1 (55)
Solano	1,497	7.5 (36)	9.9 (39)	6,388	5.2(41)	8.2(51)	13.3(20)	2,284	16.1(25)	17.6(34)		2,873	9.0 (5)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2004 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Page: 6
Run Date: 10 / 11 / 05

Comparison of Growth and Anemia Indicators by County (3,4)											
County	Birthweight (5)			Height and Weight						Anemia Low Hb/Hct (8)	
	Low		High	Short Stature (6)			Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)		
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- <95th (Rank)	% ≥95th (Rank)	Number % (Rank)
Sonoma	1,432	5.9 (12)	10.1 (43)	7,730	3.0 (2)	7.9(50)	12.1(15)	2,742	16.3(31)	16.2(21)	3,186 18.6 (56)
Stanislaus	3,622	11.5 (55)	7.8 (14)	15,929	4.1(15)	7.3(47)	14.6(32)	6,330	16.2(27)	17.0(30)	12,512 17.6 (54)
Sutter	615	7.5 (36)	12.7 (54)	2,532	3.8 (8)	8.3(54)	11.9(10)	903	16.6(39)	13.4 (8)	1,426 16.3 (48)
Tehama	381	6.6 (20)	10.8 (46)	2,044	3.8 (8)	7.1(44)	13.5(22)	804	15.5(15)	15.0(13)	1,217 19.6 (57)
Trinity	44	* *	* *	183	7.7(63)	6.6(37)	15.3(38)	66	* *	* *	101 13.9 (32)
Tulare	4,194	7.7 (40)	8.3 (24)	23,332	6.0(51)	3.8 (3)	19.1(61)	9,519	17.4(48)	18.9(44)	15,872 19.7 (58)
Tuolumne	136	2.9 (1)	11.0 (48)	741	6.2(55)	6.7(41)	11.1 (7)	282	18.8(53)	13.1 (6)	438 12.6 (22)
Ventura	4,391	6.1 (13)	8.3 (24)	18,088	4.4(22)	4.8(17)	16.2(45)	6,779	16.5(36)	18.8(43)	12,642 11.9 (16)
Yolo	726	4.1 (2)	13.2 (55)	4,486	3.7 (7)	7.2(45)	14.0(27)	1,705	15.2(14)	19.4(47)	2,570 12.8 (23)
Yuba	421	8.6 (49)	11.4 (52)	2,141	5.3(44)	6.4(34)	14.1(28)	783	15.5(15)	16.2(21)	1,173 15.9 (46)

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2004 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged < 5 Years (2)

Page: 7
Run Date: 10 / 11 / 05

Comparison of Growth and Anemia Indicators by County (3,4)													
County	Birthweight (5)			Height and Weight							Anemia Low Hb/Hct (8)		
	Low	High		Short Stature (6)	Under- weight (7)	Over- weight (7)	≥ 2 Yrs Over- weight (7)						
	Number	% <2500 g (Rank)	% >4000 g (Rank)	Number	% <5th (Rank)	% <5th (Rank)	% ≥95th (Rank)	Number	% 85th- (Rank)	% <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)
California	176,497	7.4	8.3	866,123	5.1	5.3	16.3	337,488	16.3	17.5		503,512	14.0
Nation (Prior Year)	1,172,866	9.1	7.3	4,838,257	6.2	5.2	13.6	2,044,705	15.7	14.7		3,118,041	12.8

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Infants born during the reporting period included in the analysis.

(6) Based on 2000 CDC growth chart percentiles for length-for-age for children under 2 years of age and height-for-age for children 2 years of age and older.

(7) Based on 2000 CDC growth chart percentiles for weight-for-length for children under 2 years of age and BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(8) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2004 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged 5 to <20 Years (2)

Page: 1
Run Date: 10 / 11 / 05

Comparison of Growth and Anemia Indicators by County (3,4)								
County	Height and Weight					Anemia Low Hb/Hct (7)		One or More Low/High Health Indicators (8)
	Short Stature (5)	Under- weight (6)	Over- weight (6)					
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)	% (Rank)
Alameda	13,613	6.9 (33)	3.0 (52)	17.9 (21)	22.2 (29)	7,040	15.0 (46)	* *
Alpine	85	* *	* *	* *	* *	87	* *	* *
Amador	396	6.8 (31)	3.0 (52)	16.7 (8)	23.5 (40)	298	13.1 (33)	* *
Butte	2,886	8.2 (51)	2.7 (42)	15.9 (4)	19.6 (13)	2,695	11.9 (20)	* *
Calaveras	566	6.0 (14)	1.6 (8)	14.5 (1)	23.0 (33)	528	14.8 (44)	* *
City Of Be	50	* *	* *	* *	* *	42	* *	* *
Colusa	506	7.5 (42)	2.4 (30)	18.4 (30)	23.7 (44)	458	9.6 (7)	* *
Contra Cos	6,179	6.6 (27)	2.0 (17)	18.7 (37)	24.2 (48)	3,297	14.1 (41)	* *
Del Norte	383	6.3 (23)	1.6 (8)	19.6 (47)	24.3 (50)	329	9.7 (8)	* *
El Dorado	889	7.0 (34)	2.5 (39)	17.3 (12)	16.3 (1)	712	7.4 (2)	* *
Fresno	32,682	7.8 (49)	2.4 (30)	17.3 (12)	23.1 (35)	28,008	11.6 (15)	* *
Glenn	331	6.6 (27)	1.8 (13)	23.3 (58)	19.9 (14)	309	17.5 (56)	* *

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

(8) One or More Low/High Health Indicators includes percent of children with one or more of these abnormal health indicators: short stature, underweight, at risk of overweight, overweight, or anemia. Children with more than one low/high health indicators are counted only once in the total percent. If any one expected growth or anemia indicator is missing, the percent of children with one or more low/high health indicators is not calculated.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2004 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged 5 to <20 Years (2)

Page: 2
Run Date: 10 / 11 / 05

Comparison of Growth and Anemia Indicators by County (3,4)								
County	Height and Weight					Anemia Low Hb/Hct (7)		One or More Low/High Health Indicators (8)
	Short Stature (5)	Under- weight (6)	Over- weight (6)					
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)	% (Rank)
Humboldt	1,018	4.9 (5)	1.6 (8)	18.5 (34)	24.8 (58)	762	13.5 (37)	* *
Imperial	2,424	4.8 (4)	2.8 (48)	18.3 (27)	24.3 (50)	2,436	6.4 (1)	* *
Inyo	218	5.0 (7)	1.4 (6)	25.7 (60)	21.1 (22)	143	14.0 (39)	* *
Kern	15,886	5.7 (12)	2.9 (49)	17.6 (16)	23.5 (40)	13,807	11.5 (14)	* *
Kings	1,739	6.1 (16)	2.3 (26)	19.9 (48)	23.5 (40)	1,656	21.2 (60)	* *
L.A. East	26,495	6.0 (14)	2.9 (49)	18.3 (27)	21.1 (22)	26,727	13.6 (38)	* *
L.A. North	21,995	7.5 (42)	2.3 (26)	19.0 (41)	19.9 (14)	22,415	9.0 (4)	* *
L.A. Other	97,207	4.5 (2)	2.4 (30)	18.7 (37)	25.4 (59)	75,728	12.6 (25)	* *
L.A. South	39,746	6.2 (20)	2.7 (42)	19.4 (44)	17.9 (6)	41,467	9.8 (9)	* *
L.A. West	6,315	6.1 (16)	2.2 (23)	18.7 (37)	22.6 (31)	5,839	17.4 (55)	* *
Lake	512	5.5 (11)	2.1 (21)	20.3 (49)	19.9 (14)	442	13.3 (36)	* *
Lassen	348	14.7 (60)	2.9 (49)	14.7 (2)	24.4 (52)	284	12.3 (23)	* *

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

(8) One or More Low/High Health Indicators includes percent of children with one or more of these abnormal health indicators: short stature, underweight, at risk of overweight, overweight, or anemia. Children with more than one low/high health indicators are counted only once in the total percent. If any one expected growth or anemia indicator is missing, the percent of children with one or more low/high health indicators is not calculated.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2004 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged 5 to <20 Years (2)

Page: 3
Run Date: 10 / 11 / 05

Comparison of Growth and Anemia Indicators by County (3,4)								
County	Height and Weight					Anemia Low Hb/Hct (7)		One or More Low/High Health Indicators (8)
	Short Stature (5)	Under- weight (6)	Over- weight (6)					
	Number (Rank)	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number (Rank)	% (Rank)	% (Rank)
Long Beach	6,493	4.7 (3)	2.4 (30)	19.5 (46)	23.9 (46)	5,144	14.3 (42)	* *
Madera	3,124	6.7 (30)	2.0 (17)	18.4 (30)	22.7 (32)	3,236	16.2 (51)	* *
Marin	804	9.1 (54)	2.5 (39)	20.4 (50)	24.5 (53)	533	7.7 (3)	* *
Mariposa	123	4.1 (1)	2.4 (30)	25.2 (59)	17.9 (6)	117	9.4 (6)	* *
Mendocino	1,042	7.0 (34)	1.4 (6)	22.3 (56)	20.4 (18)	687	13.0 (32)	* *
Merced	4,310	9.3 (56)	2.2 (23)	17.8 (19)	23.7 (44)	4,408	14.9 (45)	* *
Modoc	99	* *	* *	* *	* *	92	* *	* *
Mono	93	* *	* *	* *	* *	93	* *	* *
Monterey	3,513	7.5 (42)	1.3 (4)	19.1 (42)	24.5 (53)	3,381	10.7 (13)	* *
Napa	1,239	6.1 (16)	2.3 (26)	19.4 (44)	24.0 (47)	761	17.3 (54)	* *
Nevada	393	6.6 (27)	1.8 (13)	16.5 (7)	16.8 (2)	256	9.0 (4)	* *
Orange	22,946	11.1 (58)	4.1 (58)	18.3 (27)	23.3 (37)	29,444	12.7 (27)	* *

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

(8) One or More Low/High Health Indicators includes percent of children with one or more of these abnormal health indicators: short stature, underweight, at risk of overweight, overweight, or anemia. Children with more than one low/high health indicators are counted only once in the total percent. If any one expected growth or anemia indicator is missing, the percent of children with one or more low/high health indicators is not calculated.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2004 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged 5 to <20 Years (2)

Page: 4
Run Date: 10 / 11 / 05

Comparison of Growth and Anemia Indicators by County (3,4)								
County	Height and Weight					Anemia Low Hb/Hct (7)		One or More Low/High Health Indicators (8)
	Short Stature (5)	Under- weight (6)	Over- weight (6)					
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)	% (Rank)
Pasadena	1,315	4.9 (5)	3.0 (52)	17.5 (14)	20.9 (20)	1,221	10.1 (11)	* *
Placer	1,092	8.2 (51)	5.2 (60)	16.8 (10)	19.4 (10)	868	12.9 (29)	* *
Plumas	206	9.2 (55)	1.9 (16)	17.5 (14)	19.4 (10)	155	11.6 (15)	* *
Riverside	18,801	6.2 (20)	2.6 (41)	17.8 (19)	20.9 (20)	16,962	12.9 (29)	* *
Sacramento	23,532	7.6 (45)	2.7 (42)	16.7 (8)	20.1 (17)	17,010	17.5 (56)	* *
San Benito	537	8.0 (50)	1.3 (4)	20.5 (51)	23.3 (37)	558	11.8 (18)	* *
San Bernar	26,514	5.9 (13)	2.7 (42)	17.2 (11)	21.2 (24)	24,141	15.2 (47)	* *
San Diego	27,864	6.3 (23)	2.7 (42)	18.2 (25)	21.7 (27)	25,707	12.1 (21)	* *
San Franci	4,627	7.6 (45)	3.2 (57)	17.7 (18)	19.2 (9)	1,941	11.6 (15)	* *
San Joaqui	10,297	7.2 (40)	3.1 (56)	17.6 (16)	20.7 (19)	4,793	14.4 (43)	* *
San Luis O	1,470	6.8 (31)	2.7 (42)	18.0 (22)	21.6 (26)	930	12.9 (29)	* *
San Mateo	3,914	7.1 (39)	1.6 (8)	20.6 (52)	25.7 (60)	2,341	10.0 (10)	* *

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

(8) One or More Low/High Health Indicators includes percent of children with one or more of these abnormal health indicators: short stature, underweight, at risk of overweight, overweight, or anemia. Children with more than one low/high health indicators are counted only once in the total percent. If any one expected growth or anemia indicator is missing, the percent of children with one or more low/high health indicators is not calculated.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2004 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged 5 to <20 Years (2)

Page: 5
Run Date: 10 / 11 / 05

Comparison of Growth and Anemia Indicators by County (3,4)									
County	Height and Weight					Anemia Low Hb/Hct (7)		One or More Low/High Health Indicators (8)	
	Short Stature (5)		Under- weight (6)	Over- weight (6)					
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)	% (Rank)	
Santa Barb	5,288	7.0 (34)	1.7 (12)	20.6 (52)	24.5 (53)	5,028	13.2 (35)	*	*
Santa Clar	12,772	7.6 (45)	3.0 (52)	18.0 (22)	21.8 (28)	6,394	12.4 (24)	*	*
Santa Cruz	1,810	7.7 (48)	1.1 (2)	20.8 (54)	23.1 (35)	1,547	11.8 (18)	*	*
Shasta	1,681	6.1 (16)	1.8 (13)	18.2 (25)	17.1 (3)	1,156	13.1 (33)	*	*
Sierra	23	* *	* *	* *	* *	15	* *	*	*
Siskiyou	579	6.2 (20)	1.2 (3)	20.9 (55)	17.8 (5)	505	15.2 (47)	*	*
Solano	2,559	7.0 (34)	2.3 (26)	19.3 (43)	23.5 (40)	1,735	14.0 (39)	*	*
Sonoma	2,819	6.4 (25)	2.4 (30)	18.8 (40)	23.0 (33)	1,507	15.2 (47)	*	*
Stanislaus	7,383	7.0 (34)	2.2 (23)	18.1 (24)	24.2 (48)	7,588	16.7 (52)	*	*
Sutter	815	6.4 (25)	4.3 (59)	16.3 (5)	19.0 (8)	682	16.7 (52)	*	*
Tehama	758	5.1 (8)	2.4 (30)	18.5 (34)	19.4 (10)	659	17.9 (58)	*	*
Trinity	126	11.1 (58)	0.0 (1)	23.0 (57)	24.6 (56)	100	19.0 (59)	*	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

(8) One or More Low/High Health Indicators includes percent of children with one or more of these abnormal health indicators: short stature, underweight, at risk of overweight, overweight, or anemia. Children with more than one low/high health indicators are counted only once in the total percent. If any one expected growth or anemia indicator is missing, the percent of children with one or more low/high health indicators is not calculated.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.

Table 6B
2004 Pediatric Nutrition Surveillance (1)
California
Comparison of Growth and Anemia Indicators by County
Children Aged 5 to <20 Years (2)

Page: 6
Run Date: 10 / 11 / 05

Comparison of Growth and Anemia Indicators by County (3,4)								
County	Height and Weight					Anemia Low Hb/Hct (7)		One or More Low/High Health Indicators (8)
	Short Stature (5)	Under- weight (6)	Over- weight (6)					
	Number	% <5th (Rank)	% <5th (Rank)	% 85th- <95th (Rank)	% ≥95th (Rank)	Number	% (Rank)	% (Rank)
Tulare	11,054	7.3 (41)	2.4 (30)	18.6 (36)	23.3 (37)	10,621	15.2 (47)	* *
Tuolumne	404	8.9 (53)	2.0 (17)	16.3 (5)	21.5 (25)	352	12.2 (22)	* *
Ventura	6,394	5.1 (8)	2.4 (30)	18.4 (30)	24.7 (57)	6,278	10.2 (12)	* *
Yolo	2,230	5.1 (8)	2.0 (17)	18.4 (30)	22.4 (30)	1,543	12.6 (25)	* *
Yuba	808	10.8 (57)	2.1 (21)	15.2 (3)	17.1 (3)	704	12.8 (28)	* *
California	494,440	6.5	2.6	18.3	22.4	426,825	12.7	*

(1) Reporting period is January 1 through December 31.

(2) Analyses based on one record per child.

(3) Excludes records with unknown data or errors.

(4) Rank compares this county's rate to other counties. Rank 1 = best rate.

(5) Based on 2000 CDC growth chart percentiles for height-for-age for children 2 years of age and older.

(6) Based on CDC growth chart percentiles for BMI-for-age for children 2 years of age and older. 85th - < 95th percentile category identifies children at risk of overweight.

(7) Based on 1998 CDC MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States", altitude adjusted, children 6 months of age and older included in the analysis.

(8) One or More Low/High Health Indicators includes percent of children with one or more of these abnormal health indicators: short stature, underweight, at risk of overweight, overweight, or anemia. Children with more than one low/high health indicators are counted only once in the total percent. If any one expected growth or anemia indicator is missing, the percent of children with one or more low/high health indicators is not calculated.

* Percentages and ranks are not calculated if < 100 records are available for analysis after exclusions.